

A remark about unitary representations of Lorentz group and an equation for zero mass particles.

N.B. Skachkov

*Joint Institute for Nuclear Research, Laboratory of Nuclear Problems,
JINR, Dubna, 141980, RUSSIA*

† *E-mail: skachkov@cv.jinr.ru*

Abstract

In the present note the expansion of the wave function of a massless particle (with the definite value of its helicity) over the unitary irreducible representations of the Lorentz group (defined on the light cone) is used as for the analog of the Fourier transformation for deriving of an equation in the relativistic configuration representation.

Key-words: relativistic equations, Lorentz group, spin, massless particles.

The functions that realize unitary irreducible (infinite dimensional) representations of the Lorentz group [1, 2] have found their application in particle physics for formulation of relativistic equations in the so called "relativistic configurational" representation introduced in [3]. The functions used in [3] compose a complete and orthonormal set on a three dimensional mass shell hyperboloid surface ($p^2 = p_0^2 - \mathbf{p}^2 = m^2$) in Minkovsky 4-momenta space. They were found out in [2] and have a form:

$$\xi(\mathbf{p}, \mathbf{r}) = \left(\frac{p_0 - \mathbf{p}\mathbf{n}}{m} \right)^{-1-irm} \quad (1)$$

Here the $\mathbf{n}$ is a unit vector and m is the mass of a particle (in what follows we shall put $m=c=\hbar=1$). The parameter r enters the expressions for the eigenvalues of two Casimir operators of Lorentz group that are (in a general case of nonzero spin particles, i.e. for nonzero values of ν) given by formulae:

$$\hat{F} = 1/2 M_{\mu\nu} M^{\mu\nu} = \vec{M}^2 - \vec{N}^2 \implies (r^2 + \nu^2) \quad (2)$$

$$\hat{G} = 1/4 \epsilon_{\mu\nu\alpha\beta} M^{\mu\nu} M^{\alpha\beta} = \vec{M} \vec{N} \implies (r * \nu) \quad (3)$$

In [3] it was shown that spin zero functions $\xi(\mathbf{p}, \mathbf{r})$ (they correspond to a case of $\nu = 0$) do satisfy an equation

$$\hat{H}_0 \xi(\vec{p}, \vec{r}) = 2E_p \xi(\vec{p}, \vec{r}) \quad (4)$$

where the Hamiltonian $\hat{H}_0$ has a form of finite difference operator

$$\hat{H}_0 = 2\cosh(i\frac{\partial}{\partial r}) + \frac{2i}{r}\sinh(i\frac{\partial}{\partial r}) + \frac{\vec{L}^2}{(r)^2}e^{-i\frac{\partial}{\partial r}}. \quad (5)$$

The functions that realize the unitary irreducible representations of the Lorentz group in a case of non zero spin were found for a case of mass shell hyperboloid in papers [2, 4, 5]. The formulae relevant to the case of helicity basis were derived in [6]. The first attempt to write an equation analogous to (4) and (5) for the spin states [7] on a mass shell hyperboloid $p^2 = m^2$ was not successful as it was based on the functions found in [8, 9], where some phase factors were omitted. The form of a free Hamiltonian in a case of a wave function for spin 1/2 state was derived later in [10] and then extended to a case of spin 1 states [11, 12, 13]. The inclusion of interaction in this equation was done in [14]. Here a case of mass zero states (i.e. with the 4-momentum p belonging to the light cone $p^2 = p_0^2 - \mathbf{p}^2 = 0$) would be discussed. The expansion of the wave function over the complete set of the functions that realize the unitary irreducible representations on the cone manifold has the following form [15]:

$$\Psi(\mathbf{p}, \sigma) = \int_{-\infty}^{\infty} dr \int d\Omega(n) p^{-1+ir} \delta_{\sigma\nu} \delta(\mathbf{n} - \mathbf{n}_{\mathbf{p}}) \phi_{\mathbf{r}, \nu, \mathbf{n}}, \quad (6)$$

$$\phi_{\rho, \nu, \mathbf{n}} = \frac{1}{4\pi} \int \frac{d^3\mathbf{p}}{|\mathbf{p}|} p^{-1-ir} \delta_{\nu\sigma} \delta(\mathbf{n} - \mathbf{n}_{\mathbf{p}}) \Psi(\mathbf{p}, \nu). \quad (7)$$

The function

$$\zeta_{\nu}^{\sigma}(\mathbf{p}, \mathbf{r}) = p^{-1+ir} \delta_{\nu\sigma} \delta(\mathbf{n} - \mathbf{n}_{\mathbf{p}}) \quad (8)$$

can be treated as the analog of the “massive” plane waves (1). Due to the presence of the δ -functions $\delta(\mathbf{n} - \mathbf{n}_{\mathbf{p}})$ under the integrals the transformation from the momentum to r -space has in fact a one-dimensional form. The “radial” part of this plane wave

$$R(p, r) = p^{-1+ir} \quad (9)$$

obeys the following equation

$$\hat{H}_0 R(p, r) = p * R(p, r) \quad (10)$$

where the Hamiltonian has a form:

$$\hat{H}_0 = e^{-i\frac{\partial}{\partial r}}. \quad (11)$$

The connection of the transformation (6), (7) with that one based on the function (1) and (which was used in [3]) can be easily seen, for example, from the formulae of the expansion for $(-r, -\nu)$ representation given in [15]

$$\Psi(\mathbf{p}, \nu) = \int_0^\infty dr \int d\Omega(\mathbf{n}) \mathbf{p}^{-1+ir} \bar{A}_{r\nu} [1 - (\mathbf{n}\mathbf{n}_\mathbf{p})]^{-1+ir} \bar{Q}_\nu(\mathbf{n}, \mathbf{n}_\mathbf{p}) \phi_{-r, -\nu, \mathbf{n}}. \quad (12)$$

$$\phi_{-r, -\nu, \mathbf{n}} = \frac{1}{4\pi} \int \frac{d^3\mathbf{p}}{|\mathbf{p}|} p^{-1-ir} A_{r\nu} [1 - (\mathbf{n}\mathbf{n}_\mathbf{p})]^{-1-ir} Q_\nu(\mathbf{n}, \mathbf{n}_\mathbf{p}) \Psi(\mathbf{p}, \nu). \quad (13)$$

$$(14)$$

Here the coefficients $A_{r\nu}$ and $Q_\nu(\mathbf{n}, \mathbf{n}_\mathbf{p})$ are given as

$$|A_{r\nu}|^2 = \frac{1}{(4\pi)^2} (r^2 + \nu^2) \quad (15)$$

$$Q_\nu(\mathbf{n}, \mathbf{n}_\mathbf{p}) = e^{-i\nu 0(n, n_p)} (-1)^{s+\nu} \quad (16)$$

and the following relation from [15]

$$\frac{r}{(4\pi)^2} \int [1 - (\mathbf{l}\mathbf{n})]^{-1+ir} [1 - (\mathbf{l}\mathbf{k})]^{-1-ir} d\Omega(\mathbf{l}) = \delta(\mathbf{n} - \mathbf{k}). \quad (17)$$

are usefull.

References

- [1] M.A.Naimark, “Linear representations of the Lorentz group” (Pergamon Press, London, 1964).
- [2] I.S.Shapiro, Sov.Phys.Doklady, 4 (1956) 647.
- [3] V.G.Kadyshevsky, R.M.Mir-Kasimov and N.B.Skachkov, Nuovo Cim. A55 (1968) 233; Sov. J. Part. Nucl., 2 (1973) 69.
- [4] Chou Kuang-Chao and L.G.Zastavenko, Sov.J.JETP, 35(8) (1959) 990.
- [5] V.S.Popov, Sov.Phys.JETP, 10 (1960) 794.
- [6] S.Sht.Mavrodiev, N.B.Skachkov, Sov.J.Theor.Math.Phys. 23 (1975) 32.
- [7] M.D.Mateev, R.M.Mir-Kasimov and N.B.Skachkov, Sov.J.Theor.Math.Phys. 10 (1972) 3.
- [8] M.A.Liberman et.al, Sov.J.Nucl.Phys. 7 (1968) 202.
- [9] G.I.Kuznetsov et.al, Sov.J.Nucl.Phys. 10 (1969) 641.
- [10] R.A.Frick (Frik), JETP Lett. 39 (1984) 89.

- [11] R.A.Frick, Sov.J.Nucl.Phys. 38 (1983) 481.
- [12] R.A.Frick, J.Math.Phys. 38 (1997) 3457.
- [13] R.A.Frick, Eur.Phys.J. C22 (2001) 581.
- [14] R.A.Frick and N.B.Skachkov, Preprint JINR E-2-91-449, JINR, Dubna (1991).
- [15] L.G.Zastavenko and Chou Kuang-Chao , Sov.J.JETP, 11 (1960) 97.